

CANNA CLASSIC:

GROWING IN RUN-TO-WASTE



CANNA CLASSIC VEGA

The growing phase of the plant is most important. In this period the plant lays the basis for the flowering phase and a high yield. The nutrients in CANNA Classic Vega contain all the essential elements for optimal growing during this phase.

Advantages of CANNA Classic Vega

- Directly absorbable nitrogen compounds and trace elements
- Suitable for all inert substrates
- Dissolves directly
- Vital side shoots and luxuriant root development



CANNA CLASSIC FLORES

In the blooming phase the plant has different nutrient needs. During this period less nitrogen is required and there is an increased need for phosphorous and potassium. Classic Flores is rich in these elements and the special chelated trace elements allow direct absorption resulting in a perfect bloom.

Advantages of CANNA Classic Flores

- Contains all the essential ingredients for lush flowering and yield
- Classic Flores stimulates fructification and provides characteristic flavour
- Dissolves directly





CANNA RHIZOTONIC

CANNA RHIZOTONIC is a powerful, algae based, vegetative stimulator for plant roots. It contains multiple vitamins and is 100% natural. The more than 60 microbiological substances enhance the plant's hormone production and overall plant arowth.

Besides stimulating root growth on plants that are already rooted, RHIZOTONIC also stimulates new root growth on cuttings and transplants. The various trace elements and vitamins such as B1&B2 stimulate the production of hormones and root growth, thus providing beautiful, strong white roots.



CANNAZYM

CANNAZYM is a high-quality enzyme product that speeds up the process of breaking down dead root material and activates beneficial microorganisms. The dead roots are turned into sugars, which make up a valuable source of nutrients for the plant.

CANNAZYM consists of different enzymes to which vitamins and extracts are added. This improves the plant's defence system and also stimulates the plant to form new roots.



CANNABOOST ACCELERATOR

CANNABOOST Accelerator boosts the metabolism of your plants. This is an important factor, because the uptake of nutrients depends greatly on the plant's health and metabolism rate.

This additive stimulates the development of new flowers.

CANNABOOST isn't a nutrient but an additive that increases the plant's photosynthesis. This ensures that the fruits form faster and give them a better taste.



CANNA PK 13/14

CANNA PK 13/14 is a mixture of top quality nutritional minerals that stimulate flowering. It's easy- to-use and makes high yields. PK 13/14 is suitable for any growing medium.

Furthermore, it only needs to be applied for one week to the nutrient reservoir. If you do this at the right moment, you will get astonishing results in the bloom phase. Make sure to read the directions of use for the best benefits.



GROW SCHEDULE CLASSIC



Cultivation **period** in weeks

Light / Day

Substra Vega ml B/10 litres ml A/10 litres

Substra Flores ml B/10 litres ml A/10 litres

> RHIZOTONIC ml/10 litres

> > CANNAZYM CANNABOOST

PK 13/14

in mS/cm

EC Total

ml/10 litres



































4		-	ì
		· 왕 13/1	
	1		







果。
A SA



ਨ +

VEGETATIVE PHASE

Water the substrate Start / rooting (3 - 5 days)

Vegetative phase I -Plants develop in volume

0-3

8

15 - 25

20

25

0.9 - 1.3

1.3 - 1.7

0.7 - 1.1

1.1 - 1.5

^

8

10 - 20

40

fructification or appearance of the formation of flowers Vegetative phase II - Up to growth stagnation after

2 - 4

12

20 - 30

20

25

20

.2 - 1.6

1.6 - 2.0

Growth in height achieved Generative Period I - Flowers or fruits develop in length

2-3

12

25

- 35

C

25

20 - 40

1.4 - 1.8

1.8 - 2.2

Generative period II

FLOWERING

Development of the volume (breadth) of flowers or fruit

Generative Period III -

Development of the mass (weight) of flowers or fruit

2 - 3

12

15

- 25

G

25

20

- 40

1.0 - 1.4

1.4 - 1.8

25

- 50

20

- 40

0.0

0.4

12

25

-35

Cī

25

20

6

15

1.5 - 1.9

1.9 - 2.3

GENERATIVE PHASE

1 - 2

10-12

Generative Period IV -

Flowers or fruit ripening process

⋜	₹
ō	ાં.
≠	7
ਜ	ĕ
- *	≝.
<u>0</u>	Q
$\overline{\alpha}$	σ
⊐	≤
ਨ	윽
7	<u>e</u> .
ř	Š
\geq	Q
≝ ∙	<u>o</u>
⊒.	Š
3	4
±	ನ
덣.	₹
Ξ,	Q
ᆠ	\circ
ದ	ĭ
S	=
Ψ	\mathbf{Z}
⋤	W
= = .	ರ
Ξ	ŏ
₹	Ω.
Φ	<u>a</u> :
Φ	S
\supset	₽
σ	ヹ
6	=
	7
	5
2	궂
\exists	Ä
ನ	*
Mother plants remain in this phase until the end (6 - 12 months).	O.
寺	$\overline{\Delta}$
S	$\stackrel{\sim}{\sim}$
	멑
	; ; ;
	2
	This period varies depending on the species and number of plants pe

•

- The changeover from 18 to 12 hours varies depending on the variety The rule of thumb is to change after 2 weeks.
- Reduce hours of light if ripening goes too fast
- Double CANNAZYM dosage to 50 ml/10 litres, if substrate is reused. Watch out for increasing relative humidity.
- 9 20 ml/10 litres standard. Increase to a maximum of 40 ml/10 litres.
- for extra flowering power

EC: EC+ value is based in mS/cm when EC water = 0.0 at 25°C, pH 6.0. The EC total in the example is with tap water with an EC of 0.4. Add the EC of the tap water that is used to the recommended EC!

Recommended pH is between 5.2 and 6.2

Use pH- grow in the vegetative phase to lower the pH Use pH- bloom in the generative phase to lower the pH Adding pH- can increase EC.

> percentage in substrate, water dosage strategy, etc. humidity, plant species, root volume, moisture is further determined by factors such as: temperature fertilization strategy. The optimum fertilization strategy can help novice growers to develop a sophisticated The guidelines in the table aren't an iron law, but